

What is claimed is:

1. A method of controlling the formation of crystalline hydrates in a fluid system, said method comprising contacting the fluid with a polymer capable of interacting with charged molecules in the fluid, and allowing the polymer to sorb the charged gaseous molecules.

10

15

20

25

30

2. The method as claimed in claim 1 wherein the fluid is selected from the group consisting essentially of

- (i) oil and water,
- (ii) gas and water,
- (iii) a combination of oil and gas and water.

3. The method as claimed in claim 1 wherein the polymer is a hyperbranched polyamino polymer.

4. The method as claimed in claim 1 wherein the polymer is a dendritic polymer.

5. The method as claimed in claim 1 wherein the polymer is a combination of a

hyperbranched polyamino polymer and a dendritic polymer.

6. The method as claimed in claim 1 wherein the polymer is a siliconized hyperbranched polyamino polymer.

7. The method as claimed in claim 1 wherein the polymer is a siliconized dendritic polymer.

8. The method as claimed in claim 1 wherein the polymer is a combination of a siliconized hyperbranched polyamino polymer and a siliconized dendritic polymer.

9. The method as claimed in claim 1 wherein the polymer is a combination of a hyperbranched polyamino polymer and a dendritic polymer and wherein one of the polymers is siliconized.

10. A method as claimed in claim 1 wherein the polymer has a molecular weight of at least 5000 Daltons.

11. A method as claimed in claim 1 wherein the polymer is associated with a solid particle.

12. A method as claimed in claim 11 wherein the association is the immobilization of the polymer on the surface of the solid particle.

13. A method as claimed in claim 11 wherein the association is the solid particle embedded in the polymer.

14. A method as claimed in claim 11 wherein the solid particle is silica.

15. A method as claimed in claim 14 wherein the silica is fumed.

16. A method as claimed in claim 14 wherein the silica is precipitated.

17. A method as claimed in claim 14 wherein the silica is a silica gel.

18. A method as claimed in claim 14 wherein the silica is dispersed.
19. A method as claimed in claim 11 wherein the solid particle is diatomaceous earth.
20. A method as claimed in claim 11 wherein the solid particle is sand.
21. A method as claimed in claim 11 wherein the solid particle is cellulose.
- 5 22. A method as claimed in claim 11 wherein the solid particle is polystyrene.
23. A method as claimed in claim 11 wherein the solid particle is clay.
24. A method as claimed in claim 11 wherein the solid particle is porous.
25. A method as claimed in claim 11 wherein the solid particle is nonporous.
26. A method as claimed in claim 11 wherein the solid particle is hydrophobic.
- 10 27. A method as claimed in claim 11 wherein the solid particle is hydrophilic.
28. A method as claimed in claim 11 wherein the solid particle is a nano particle.
29. A method as claimed in claim 11 wherein the solid particle is a macro particle.
30. A method as claimed in claim 11 wherein the solid particle is a micro particle.
31. In combination, a mixture of a fluid and a polymer capable of interacting with
15 charged molecules in the fluid.
32. In combination, a mixture of a fluid and a polymer capable of interacting with
charged molecules in the fluid wherein the polymer is associated with a solid particle.

20

25

30